



03

University of Sri Jayewardenepura  
Faculty of Humanities and Social Sciences  
Bachelor of Arts Third Year Second Semester Examination- 2019 January/February

**Economics****ECON 3260.03 - Econometrics**

Time: Three hours (03)

**Answer FIVE (05) questions including FIRST QUESTION.**

Calculators are allowed. Statistical tables are provided.

Each question carries equal marks.

01. Briefly explain the following terms.

- I. Goodness of fit
- II. Type I error
- III. Correlation coefficient
- IV.  $\beta$  coefficients
- V. Overall significance of a regression
- VI. Linear regression
- VII. Tolerance
- VIII. First order auto regressive scheme

(02 marks each)

02. Explain the steps of an econometric research in terms of your own example.

(16 marks)

03. Consider the following intermediate results.

$$\begin{array}{lll} \sum y = 135 & \sum x_1 = 105 & \sum x_2 = 180 \\ \bar{y} = 9 & \bar{x}_1 = 7 & \bar{x}_2 = 12 \\ \sum x_1 y = -28 & \sum x_1^2 = 60 & \sum y^2 = 40 \\ \sum x_2 y = 38 & \sum x_2^2 = 24 & \sum \hat{y}^2 = 27.73 \\ \sum x_1 x_2 = -12 & n = 15 & \end{array}$$

- I. Estimate least squares regression equation and interpret. (04 marks)
- II. Test the statistical significance of the regression coefficients  $\hat{\beta}_1$  and  $\hat{\beta}_2$  at 5% level of significance. (06 marks)
- III. Evaluate the goodness of fit of the model. (02 marks)
- IV. Construct confidence intervals for slope coefficients  $\hat{\beta}_1$  and  $\hat{\beta}_2$  at 5% level of significance and interpret. (04 marks)

04. I. Briefly explain the stochastic assumptions of ordinary least squares method. (08 marks)

II. Distinguish between each of the following pairs of concepts.

- a. Unbiasness and consistency
- b. Consistency and asymptotic efficiency
- c. Efficiency and minimum variance
- d. Unbiasness and MSE

(02 marks each)

05. The following table includes annual consumption and disposable income in a hypothetical economy (Rs mn).

$C_t$	297	303	308	325	339	338	358	369	378	391	413	432
$Y_t$	331	373	338	360	378	375	398	410	417	445	462	486
$e_t$	-3.3	1.9	2.1	0.46	0.08	0.98	1.62	-8.64	5.75	-4.98	2.29	1.71

Estimated model:  $\hat{C}_t = 21.5 + 0.841 Y_d$

(9.4) (0.024)

I. Use graphical method to identify the econometric problem of the above model.

(04 marks)

II. Conduct a formal test to detect the problem considering the additional information given below.

$$\sum e_1^2 = 68.3 \quad \sum e_2^2 = 9.6 \quad c = 3 \quad d.f = \left[ \frac{n-c}{2} \right] - K$$

(05 marks)

III. How do you remove the problematic situation of the above model? Illustrate with an example.

(04 marks)

IV. Briefly explain one alternative test which can be used to detect the above problem.

(03 marks)

06. I. What is multicollinearity? (02 marks)

II. Discuss the causes consequences of multicollinearity. (08 marks)

III. How do you conduct "Frisch's confluence analysis" to detect multicollinearity?


(03 marks)

IV. Briefly explain your suggestions to minimize the effect of multicollinearity of a model.

(03 marks)

\*\*\*\*\*

Table 1. Areas under the Normal Curve




Example

$$Z = \frac{X - \mu}{\sigma}$$

$$P(Z > 1.96) = .0250$$

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0022	.0021	.0020	.0019
2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010

Table 2. Percentage Points of the  $t$  Distribution



Example

For  $\nu = 10$  degrees of freedom:

$$P(t > 1.812) = 0.05$$

$$P(t < -1.812) = 0.05$$

$\alpha \backslash \nu$	.25	.20	.15	.10	.05	.025	.01	.005	.0005
1	1.000	1.376	1.963	3.078	6.314	12.706	31.821	63.657	636.619
2	.816	1.061	1.386	1.886	2.920	4.303	6.965	9.925	31.598
3	.765	.978	1.250	1.638	2.353	3.182	4.541	5.841	12.941
4	.744	.941	1.190	1.533	2.132	2.776	3.747	4.604	8.610
5	.727	.920	1.156	1.476	2.015	2.571	3.365	4.032	6.859
6	.718	.906	1.134	1.440	1.943	2.447	3.143	3.707	5.959
7	.711	.896	1.119	1.415	1.895	2.365	2.998	3.499	5.405
8	.706	.889	1.108	1.397	1.860	2.306	2.896	3.355	5.041
9	.703	.883	1.100	1.383	1.833	2.262	2.821	3.250	4.781
10	.700	.879	1.093	1.372	1.812	2.228	2.764	3.169	4.587
11	.697	.876	1.088	1.363	1.796	2.201	2.718	3.106	4.437
12	.695	.873	1.083	1.356	1.782	2.179	2.681	3.055	4.318
13	.694	.870	1.079	1.350	1.771	2.160	2.650	3.012	4.221
14	.692	.868	1.076	1.345	1.761	2.145	2.624	2.977	4.140
15	.691	.866	1.074	1.341	1.753	2.131	2.602	2.947	4.073
16	.690	.865	1.071	1.337	1.746	2.120	2.583	2.921	4.015
17	.689	.863	1.069	1.333	1.740	2.110	2.567	2.898	3.965
18	.688	.862	1.067	1.330	1.734	2.101	2.552	2.878	3.922
19	.688	.861	1.066	1.328	1.729	2.093	2.539	2.861	3.883
20	.687	.860	1.064	1.325	1.725	2.086	2.528	2.845	3.850
21	.686	.859	1.063	1.323	1.721	2.080	2.518	2.831	3.819
22	.686	.858	1.061	1.321	1.717	2.074	2.508	2.819	3.792
23	.685	.858	1.060	1.319	1.714	2.069	2.500	2.807	3.767
24	.685	.857	1.059	1.318	1.711	2.064	2.492	2.797	3.745
25	.684	.856	1.058	1.316	1.708	2.060	2.485	2.787	3.725
26	.684	.856	1.058	1.315	1.706	2.056	2.479	2.779	3.707
27	.684	.855	1.057	1.314	1.703	2.052	2.473	2.771	3.690
28	.683	.855	1.056	1.313	1.701	2.048	2.467	2.763	3.674
29	.683	.854	1.055	1.311	1.699	2.045	2.462	2.756	3.659
30	.683	.854	1.055	1.310	1.697	2.042	2.457	2.750	3.646
40	.681	.851	1.050	1.303	1.684	2.021	2.423	2.704	3.551
60	.679	.848	1.046	1.296	1.671	2.000	2.390	2.660	3.460
120	.677	.845	1.041	1.289	1.658	1.980	2.358	2.617	3.373
$\infty$	.674	.842	1.036	1.282	1.645	1.960	2.326	2.576	3.291



Table 4A. Values of  $F_{\alpha, \nu_1, \nu_2}$

$\nu_1$  = degrees of freedom for numerator

Example:  $\nu_1 = 9, \nu_2 = 12$  degrees of freedom  
 $P(F > 2.80) = 0.05$

$\nu_2 \backslash \nu_1$	1	2	3	4	5	6	7	8	9	10	12	15	20	24	30	40	60	120	$\infty$
1	161	200	216	225	230	234	237	239	241	242	244	246	248	249	250	251	252	253	254
2	18.5	19.0	19.2	19.2	19.3	19.3	19.4	19.4	19.4	19.4	19.4	19.4	19.4	19.4	19.5	19.5	19.5	19.5	19.5
3	10-1	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.70	8.66	8.64	8.62	8.59	8.57	8.55	8.53
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.86	5.80	5.77	5.75	5.72	5.69	5.66	5.63
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.68	4.62	4.56	4.53	4.50	4.46	4.43	4.40	4.37
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.00	3.94	3.87	3.84	3.81	3.77	3.74	3.70	3.67
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.57	3.51	3.44	3.41	3.38	3.34	3.30	3.27	3.23
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.28	3.22	3.15	3.12	3.08	3.04	3.00	2.97	2.93
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.07	3.01	2.94	2.90	2.86	2.83	2.79	2.75	2.71
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.91	2.85	2.77	2.74	2.70	2.66	2.62	2.58	2.54
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.79	2.72	2.65	2.61	2.57	2.53	2.49	2.45	2.40
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.69	2.62	2.54	2.51	2.47	2.43	2.38	2.34	2.30
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.60	2.53	2.46	2.42	2.38	2.34	2.30	2.25	2.21
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.53	2.46	2.39	2.35	2.31	2.27	2.22	2.18	2.13
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.48	2.40	2.33	2.29	2.25	2.20	2.16	2.11	2.07
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	2.49	2.42	2.35	2.28	2.24	2.19	2.15	2.11	2.06	2.01
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	2.45	2.38	2.31	2.23	2.19	2.15	2.10	2.06	2.01	1.96
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	2.41	2.34	2.27	2.19	2.15	2.11	2.06	2.02	1.97	1.92
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	2.38	2.31	2.23	2.16	2.11	2.07	2.03	1.98	1.93	1.88
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	2.35	2.28	2.20	2.12	2.08	2.04	1.99	1.95	1.90	1.84
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37	2.32	2.25	2.18	2.10	2.05	2.01	1.96	1.92	1.87	1.81
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	2.30	2.23	2.15	2.07	2.03	1.98	1.94	1.89	1.84	1.78
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	2.27	2.20	2.13	2.05	2.01	1.96	1.91	1.86	1.81	1.76
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.18	2.11	2.03	1.98	1.94	1.89	1.84	1.79	1.73
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	2.24	2.16	2.09	2.01	1.96	1.92	1.87	1.82	1.77	1.71
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.09	2.01	1.93	1.89	1.84	1.79	1.74	1.68	1.62
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.00	1.92	1.84	1.79	1.74	1.69	1.64	1.58	1.51
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	1.99	1.92	1.84	1.75	1.70	1.65	1.59	1.53	1.47	1.39
120	3.92	3.07	2.68	2.45	2.29	2.18	2.09	2.02	1.96	1.91	1.83	1.75	1.66	1.61	1.55	1.50	1.43	1.35	1.25
$\infty$	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88	1.83	1.75	1.67	1.57	1.52	1.46	1.39	1.32	1.22	1.00

$\nu_2$  = degrees of freedom for denominator

$$\hat{\beta}_0 = \frac{\sum Y_i \sum X_i^2 - \sum X_i Y_i \sum X_i}{n \sum X_i^2 - (\sum X_i)^2}$$

$$\hat{\beta}_1 = \frac{n \sum X_i Y_i - \sum X_i \sum Y_i}{n \sum X_i^2 - (\sum X_i)^2}$$

$$\hat{\beta}_1 = \frac{(\sum x_1 y)(\sum x_2^2) - (\sum x_2 y)(\sum x_1 x_2)}{(\sum x_1^2)(\sum x_2^2) - (\sum x_1 x_2)^2}$$

$$\hat{\beta}_2 = \frac{(\sum x_2 y)(\sum x_1^2) - (\sum x_1 y)(\sum x_1 x_2)}{(\sum x_1^2)(\sum x_2^2) - (\sum x_1 x_2)^2}$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X}_1 - \hat{\beta}_2 \bar{X}_2$$

$$Y_i = \beta_0 + \beta_1 \left[ \frac{1}{X_1} \right] + \mu_i$$

$$\text{Var} \hat{\beta}_1 = \sigma_u^2 \frac{\sum x_2^2}{\sum x_1^2 x_2^2 - (\sum x_1 x_2)^2}$$

$$\text{Var} \hat{\beta}_2 = \sigma_u^2 \frac{\sum x_1^2}{\sum x_1^2 x_2^2 - (\sum x_1 x_2)^2}$$

$$\ln Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \varepsilon_i$$

$$\text{Var} \mu = \frac{\sum e^2}{n - K}$$